

EUROCRINE

Clinic Benchmark Report

Surgery of the Thyroid Gland
2023

EUROCRINE Clinic Benchmark Report on Thyroid Resections performed in 2023



Investigated Clinic

- Markera alla
- Bröst och Endokrinkirurgiska kliniken, KS Huddinge
- Carlanderska Endokrinkirurgi
- Dubbel - Kirurgiska kliniken, Lund
- Gastro Center Skåne
- Kirurgicentrum Skåne
- Kirurgiska kliniken, Borås
- Kirurgiska kliniken, Danderyd
- Kirurgiska kliniken, Eksjö
- Kirurgiska kliniken, Falun
- Kirurgiska kliniken, Gävle
- Kirurgiska kliniken, Göteborg
- Kirurgiska kliniken, Halmstad
- Kirurgiska kliniken, Helsingborg
- Kirurgiska kliniken, Jönköping
- Kirurgiska kliniken, Kalmar
- Kirurgiska kliniken, Karlskrona
- Kirurgiska kliniken, Karlstad
- Kirurgiska kliniken, Kristianstad
- Kirurgiska kliniken, KS Solna
- Kirurgiska kliniken, Kungälv
- Kirurgiska kliniken, Linköping
- Kirurgiska kliniken, Ljungby
- Kirurgiska kliniken, Luleå
- Kirurgiska kliniken, Lund
- Kirurgiska kliniken, Norrköping
- Kirurgiska kliniken, Nyköping
- Kirurgiska kliniken, Örebro
- Kirurgiska kliniken, Östersund
- Kirurgiska kliniken, Skövde
- Kirurgiska kliniken, Södersjukhuset

Benchmark Clinics

- Markera alla
- Bröst och Endokrinkirurgiska kliniken, KS Huddinge
- Carlanderska Endokrinkirurgi
- Dubbel - Kirurgiska kliniken, Lund
- Gastro Center Skåne
- Kirurgicentrum Skåne
- Kirurgiska kliniken, Borås
- Kirurgiska kliniken, Danderyd
- Kirurgiska kliniken, Eksjö
- Kirurgiska kliniken, Falun
- Kirurgiska kliniken, Gävle
- Kirurgiska kliniken, Göteborg
- Kirurgiska kliniken, Halmstad
- Kirurgiska kliniken, Helsingborg
- Kirurgiska kliniken, Jönköping
- Kirurgiska kliniken, Kalmar
- Kirurgiska kliniken, Karlskrona
- Kirurgiska kliniken, Karlstad
- Kirurgiska kliniken, Kristianstad
- Kirurgiska kliniken, KS Solna
- Kirurgiska kliniken, Kungälv
- Kirurgiska kliniken, Linköping
- Kirurgiska kliniken, Ljungby
- Kirurgiska kliniken, Luleå
- Kirurgiska kliniken, Lund
- Kirurgiska kliniken, Norrköping
- Kirurgiska kliniken, Nyköping
- Kirurgiska kliniken, Örebro
- Kirurgiska kliniken, Östersund
- Kirurgiska kliniken, Skövde
- Kirurgiska kliniken, Södersjukhuset

Investigated Year

- Markera alla
- 2012
- 2013
- 2014
- 2015
- 2016
- 2017
- 2018
- 2019
- 2020
- 2021
- 2022
- 2023
- 2024

EUROCRINE Clinic Benchmark Report on Thyroid Resections performed in 2023



Registered Thyroid Resections in Investigated Clinic

231



2023

Registered Thyroid Resections in the Other EUROCRINE Clinics of Your Country

2855



2023

Color Coding in this Report:

Investigated Clinic

Other EUROCRINE Clinics in Your Country


Thyroid

Gender and Age Distribution




Investigated Clinic

Age Distribution and Gender - Benign Disease at Histology




EUROCRINE Clinics


Age Distribution and Gender - Benign Disease at Histology



Age Distribution and Gender - Malignancy at Histology



Age Distribution and Gender - Malignancy at Histology



Thyroid

Preoperative Workup: Ultrasound



Investigated Clinic

EUROCRINE Clinics


Preoperative Ultrasound of the Thyroid Gland

Preoperative Ultrasound of the Thyroid Gland



Preoperative Ultrasound of Regional Lymph Nodes for Main Indication 'Excluding Malignancy' or 'Malignancy'

Preoperative Ultrasound of Regional Lymph Nodes for Main Indication 'Excluding Malignancy' or 'Malignancy'



FNA: Performed Under Guidance of Ultrasonography

FNA: Performed Under Guidance of Ultrasonography




Thyroid

Preoperative Workup: Ultrasound and EU-TIRADS Classification (variable introduced in 2020)




Investigated Clinic

EU-TIRADS Classification Distribution




EU-TIRADS Classification and Final Diagnosis Groups on Histopathology




EUROCRINE Clinics

EU-TIRADS Classification Distribution



EU-TIRADS Classification and Final Diagnosis Groups on Histopathology





Investigated Clinic

EUROCRINE Clinics



Bethesda Classification Distribution

Bethesda Classification Distribution



Cytology Result and Main Histopathological Diagnosis Group

Cytology Result and Main Histopathological Diagnosis Group



Thyroid

Main Indication for Surgery





Investigated Clinic

EUROCRINE Clinics



Main Indication for Surgery (Primary Operations)

Main Indication for Surgery (Primary Operations)



Main Indication and Diagnosis at Histology

Main Indication and Diagnosis at Histology



Thyroid

Main Indication for Surgery: Excluding Malignancy





Investigated Clinic

EUROCRINE Clinics



Main Indication (Excluding Malignancy) and Final Diagnosis Groups

Main Indication (Excluding Malignancy) and Final Diagnosis Groups



Main Indication (Excluding Malignancy) and Final Histopathological Diagnosis

Main Indication (Excluding Malignancy) and Final Histopathological Diagnosis




Thyroid

Main Indication for Surgery: Volume and Compression Symptoms




Investigated Clinic

Main Indication (Volume and Compression Symptoms) and Final Diagnosis Groups




Main Indication (Compression Symptoms) and Final Histopathological Diagnosis




EUROCRINE Clinics

Main Indication (Volume and Compression Symptoms) and Final Diagnosis Groups



Main Indication (Compression Symptoms) and Final Histopathological Diagnosis




Thyroid

Main Indication for Surgery: Thyrotoxicosis




Investigated Clinic

Main Indication (Volume and Compression Symptoms) and Final Diagnosis Groups




Main Indication (Compression Symptoms) and Final Histopathological Diagnosis




EUROCRINE Clinics

Main Indication (Volume and Compression Symptoms) and Final Diagnosis Groups



Main Indication (Compression Symptoms) and Final Histopathological Diagnosis



Thyroid

Main Indication for Surgery: Thyrotoxicosis





Investigated Clinic

EUROCRINE Clinics



Main Indication (Malignancy) and Final Diagnosis Groups

Main Indication (Malignancy) and Final Diagnosis Groups



Main Indication (Malignancy) and Final Histopathological Diagnosis

Main Indication (Malignancy) and Final Histopathological Diagnosis



Thyroid

Surgical Approach



Investigated Clinic

EUROCRINE Clinics

Surgical Approach

Surgical Approach



Neck Incision - Subspecification

Neck Incision - Subspecification

Neck incision	Number
Conventional neck incision (regardless of length)	227

Neck incision	Number
Conventional neck incision (regardless of length)	2731
Lateral approach	4

Extracervical Approach - Subspecification

Extracervical Approach - Subspecification

Extra cervical approach	Number
Other	1

Extra cervical approach	Number
Other	4


Thyroid

Resection Type and Intrathoracic Goiter Extension




Investigated Clinic


Resection Type



Intrathoracic Goiter Extension




Need for Sterno-/Thoracotomy




EUROCRINE Clinics


Resection Type



Intrathoracic Goiter Extension




Need for Sterno-/Thoracotomy




Investigated Clinic

Intraoperative Use of IONM



Yes 100,0%

IONM Induced Change of Strategy




Yes 0,4%

No 99,6%

EUROCRINE Clinics


Intraoperative Use of IONM



No 5,7%

Yes 94,3%

IONM Induced Change of Strategy




Yes 0,3%

No 99,7%

Type of Intraoperative Nerve Monitoring (Variable introduced 2019)


● Intermittent nerve monitoring



100,00%

Type of Intraoperative Nerve Monitoring (Variable introduced 2019)

● Continous nerve monitoring
● Intermittent nerve monitoring



99,93%


Thyroid

Operation Time (skin to skin)




Hemithyroidectomy

Hemithyroidectomy, investigated clinic (graph limited to 6 hours)




Hemithyroidectomy, EUROCRINE clinics (graph limited to 6 hours)




Thyroidectomy

Thyroidectomy, investigated clinic (graph limited to 6 hours)



Thyroidectomy, EUROCRINE clinics (graph limited to 6 hours)




Thyroid


Papillary Thyroid Cancer PTC, Initial Resection Type and Extent of Lymphadenectomy (excluding Tumorstage pT1a)



Resection Type in PTC - Investigated Clinic



Thyroidectomy and Extent of Lymphadenectomy



Unilateral Lobectomy of the Thyroid Gland and Extent of Lymphadenectomy




Thyroid

Papillary Thyroid Cancer PTC, Initial Resection Type and Extent of Lymphadenectomy (excluding Tumorstage pT1a)



Resection Type in PTC - EUROCRINE Clinics




Thyroidectomy and Extent of Lymphadenectomy

- Bilateral central lymph node dissection (PJD41)
- Bilateral central lymphnode dissection AND bilateral lateral lymphno...
- Central lymph node dissection (PJD41) and one-sided lat. Lymph nod...
- Exploration of lymph nodes incl. Biopsy (PJD10)
- Extirpation of lymph nodes (PJD41)
- None
- Unilateral central lymph node dissection (PJD41)



Unilateral Lobectomy of the Thyroid Gland and Extent of Lymphadenectomy

- Central lymph node dissection (PJD41) and one-sided lat. Lymph nod...
- Extirpation of lymph nodes (PJD41)
- None
- Unilateral central lymph node dissection (PJD41)



Thyroid

Complications: Postoperative Recurrent Laryngeal Nerve Palsy (preoperatively known paresis and resected nerves excluded)



Investigated Clinic

All Operations

Patients	Number of Postop Paresis	%	% (NAR)	Paresis at Follow-up(F2)	%	% (NAR)
228	1	0,44	0,23			


Primary Operations

Patients	Number of Postop Paresis	%	% (NAR)	Paresis at Follow-up(F2)	%	% (NAR)
195	1	0,51	0,26			

Reoperations

Patients	Number of Postop Paresis	%	% (NAR)	Paresis at Follow-up(F2)	%	% (NAR)
31						

Complications Summary - Recurrent Nerve Palsy - All Operations - Latest Available Follow-up



EUROCRINE Clinics

All Operations

Patients	Number of Postop Paresis	%	% (NAR)	Paresis at Follow-up(F2)	%	% (NAR)
2833	64	2,26	1,20	11	0,25	0,206


Primary Operations

Patients	Number of Postop Paresis	%	% (NAR)	Paresis at Follow-up(F2)	%	% (NAR)
2508	55	2,19	1,14	8	0,21	0,17

Reoperations

Patients	Number of Postop Paresis	%	% (NAR)	Paresis at Follow-up(F2)	%	% (NAR)
322	9	2,80	1,76	3	0,61	0,59

Complications Summary - Recurrent Nerve Palsy - All Operations - Latest Available Follow-up




Thyroid

Recurrent Nerve Palsy - Laryngoscopy Policy (starting Feb 20th 2020, after introduction of these variables)

Investigated Clinic

Laryngoscopy Status at (F1)



Reasons for Performing Postoperative Laryngoscopy


Reason	n	%
Routine examination as department policy (>90% investigated)	1	50,00%
Targeted examination (hoarseness, loss of signal, morphological suspected nerve injury)	1	50,00%
Total	2	100,00%

Reasons for Not Performing Postoperative Laryngoscopy

Reason	n	%
Not performed on clinical grounds (normal voice)	132	58,41%
Not performed because of normal IONM signal	87	38,50%
Not performed for other reasons	7	3,10%
Total	226	100,00%

EUROCRINE Clinics

Laryngoscopy Status at (F1)



Reasons for Performing Postoperative Laryngoscopy

Reason	n	%
Routine examination as department policy (>90% investigated)	2538	88,90%
Targeted examination (hoarseness, loss of signal, morphological suspected nerve injury)	230	8,06%
Targeted examination (hoarseness, loss of signal, morphological suspected nerve injury)	87	3,05%
Total	2855	100,00%


Reasons for Not Performing Postoperative Laryngoscopy

Reason	n	%
Not performed on clinical grounds (normal voice)	1367	55,28%
Not performed because of normal IONM signal	700	28,31%
Not performed for other reasons	378	15,29%
	28	1,13%
Total	2473	100,00%

Thyroid
Intraoperative Use of Fluorescence Imaging (Variables introduced in 2023)




Overall Use, in all Resection Types




Reimplantation of Parathyroid Tissue

Without Use of Fluorescence




With Use of Fluorescence




Parathyroid Gland on Thyroid Specimen


Without Use of Fluorescence



With Use of Fluorescence




Overall Use, in all Resection Types




Reimplantation of Parathyroid Tissue

Without Use of Fluorescence




With Use of Fluorescence




Parathyroid Gland on Thyroid Specimen

Without Use of Fluorescence




With Use of Fluorescence




Thyroid
Short and Longterm Treatment of Postoperative Parathyroid Insufficiency with Calcium (all Thyroidectomies, Primary Operations)




Thyroidectomies




Calcium po at (F1)



Calcium po at next Follow-up (F2)




Complication Summary: Calcium po at Latest Available Follow-up




Thyroidectomies




Calcium po at (F1)



Calcium po at next Follow-up (F2)



Complication Summary: Calcium po at Latest Available Follow-up



Thyroid

Short and Longterm Treatment of Postoperative Parathyroid Insufficiency with Vit D (all Thyroidectomies, Primary Operations)






Thyroidectomies

Vit D po at (F1)

Vit D po at next Follow-up (F2)

Complication Summary: Vit D po at Latest Available Follow-up






Thyreoidectomies

Vit D po at (F1)

Vit D po at next Follow-up (F2)

Complication Summary: Vit D po at Latest Available Follow-up



Thyroid

Short and Longterm PTH Course (all Thyroidectomies, Primary Operations)






Thyroidectomies

PTH at (F1)

PTH at Longterm Follow-up

Complication Summary: Vit D po at Latest Available Follow...






Thyroidectomies

PTH at (F1)

PTH at Longterm Follow-up

Complication Summary: Vit D po at Latest Available Follow...



Thyroid

Lost Parathyroid Glands on the Specimen, Postoperative Hemorrhage, Wound Infection and Hospital Stay



Investigated Clinic

EUROCRINE Clinics

Parathyroid Gland on Specimen







Postoperative Hemorrhage

Wound Infection


Parathyroid Gland on Specimen

Postoperative Hemorrhage

Wound Infection



Postoperative Hospital Stay (graph limited to max 31 days)



Postoperative Hospital Stay (graph limited to max 31 days)

